


Novel H1N1 Flu

What is H1N1 (swine flu)?

H1N1 (referred to as “swine flu” early on) is a new influenza virus causing illness in people. This new virus was first detected in people in the United States in April 2009. Other countries, including Mexico and Canada, have reported people sick with this new virus. This virus is spreading from person-to-person, probably in much the same way that regular seasonal influenza viruses spread.



Why is this new H1N1 virus sometimes called “swine flu”?

This virus was originally referred to as “swine flu” because laboratory testing showed that many of the genes in this new virus were very similar to influenza viruses that normally occur in pigs in North America. But further study has shown that this new virus is very different from what normally circulates in North American pigs. It has two genes from flu viruses that normally circulate in pigs in Europe and Asia and avian genes and human genes. Scientists call this a “quadruple reassortant” virus.

Novel H1N1 Flu in Humans

Are there human infections with this H1N1 virus in the U.S.?

Yes. Cases of human infection with this H1N1 influenza virus were first confirmed in the U.S. in Southern California and near Guadalupe County, Texas. The outbreak intensified rapidly from that time and more and more states have been reporting cases of illness from this virus. An updated case count of confirmed novel H1N1 flu infections in the United States is kept at <http://www.cdc.gov/h1n1flu/investigation.htm>. CDC and local and state health agencies are working together to investigate this situation.

Is this new H1N1 virus contagious?

CDC has determined that this new H1N1 virus is contagious and is spreading from human to human. However, at this time, it is not known how easily the virus spreads between people.

What are the signs and symptoms of this virus in people?

The symptoms of this new H1N1 flu virus in people are similar to the symptoms of seasonal flu and include fever, cough, sore throat, runny or stuffy nose, body aches, headache, chills and fatigue. A significant number of people who have been infected with this virus also have reported diarrhea and vomiting. Also, like seasonal flu, severe illnesses and death has occurred as a result of illness associated with this virus.



How severe is illness associated with this novel H1N1 flu virus?

It’s not known at this time how severe this new H1N1 flu virus will be in the general population. In seasonal flu, there are certain people that are at higher risk of serious flu-related complications. This includes people 65 years and older, children younger than five years old, pregnant women, and people of any age with certain chronic medical conditions. Early indications are that pregnancy and other previously recognized medical conditions that increase the risk of influenza-related complications, like

asthma and diabetes, also appear to be associated with increased risk of complications from this novel H1N1 virus infection as well.

One thing that appears to be different from seasonal influenza is that adults older than 64 years do not yet appear to be at increased risk of novel H1N1-related complications thus far in the outbreak. CDC is conducting laboratory studies to see if certain people might have natural immunity to this virus, depending on their age. Early reports indicate that no children and few adults younger than 60 years old have existing antibody to the novel H1N1 flu virus; however, about one-third of adults older than 60 may have antibodies against this virus. It is unknown how much, if any, protection may be afforded against the novel H1N1 flu by any existing antibody.

How does novel H1N1 flu compare to seasonal flu in terms of its severity and infection rates?

CDC is still learning about the severity of the novel H1N1 flu virus. At this time, there is not enough information to predict how severe this novel H1N1 flu outbreak will be in terms of illness and death or how it will compare with seasonal influenza.

With seasonal flu, we know that seasons vary in terms of timing, duration and severity. Seasonal influenza can cause mild to severe illness, and at times can lead to death. Each year, in the United States, on average 36,000 people die from flu-related complications and more than 200,000 people are hospitalized from flu-related causes. Of those hospitalized, 20,000 are children younger than 5 years old. Over 90% of deaths and about 60 percent of hospitalization occur in people older than 65.

So far, with novel H1N1 flu, the largest number of novel H1N1 flu confirmed and probable cases have occurred in people between the ages of 5 and 24-years-old. At this time, there are few cases and no deaths reported in people older than 64 years old, which is unusual when compared with seasonal flu. However, pregnancy and other previously recognized high risk medical conditions from seasonal influenza appear to be associated with increased risk of complications from this novel H1N1.

How does this new H1N1 virus spread?

Spread of this H1N1 virus is thought to be happening in the same way that seasonal flu spreads. Flu viruses are spread mainly from person to person through coughing or sneezing by people with influenza. Sometimes people may become infected by touching something with flu viruses on it and then touching their mouth or nose.

How long can an infected person spread this virus to others?

At the current time, CDC believes that this virus has the same properties in terms of spread as seasonal flu viruses. With seasonal flu, studies have shown that people may be contagious from one day before they develop symptoms to up to 7 days after they get sick. Children, especially younger children, might potentially be contagious for longer periods. CDC is studying the virus and its capabilities to try to learn more and will provide more information as it becomes available.

Exposures Not Thought to Spread New H1N1 Flu

Can I get infected with this new H1N1 virus from eating or preparing pork?

No. H1N1 viruses are not spread by food. You cannot get this new H1N1 virus from eating pork or pork products. Eating properly handled and cooked pork products is safe.

Is there a risk from drinking water?

Tap water that has been treated by conventional disinfection processes does not likely pose a risk for transmission of influenza viruses. Current drinking water treatment regulations provide a high degree of protection from viruses. No research has been completed on the susceptibility of the novel H1N1 flu virus to conventional drinking water treatment processes. However, recent studies have demonstrated that free chlorine levels typically used in drinking water treatment are adequate to inactivate highly pathogenic

H5N1 avian influenza. It is likely that other influenza viruses such as novel H1N1 would also be similarly inactivated by chlorination. To date, there have been no documented human cases of influenza caused by exposure to influenza-contaminated drinking water.

Can the new H1N1 flu virus be spread through water in swimming pools, spas, water parks, interactive fountains, and other treated recreational water venues?

Influenza viruses infect the human upper respiratory tract. There has never been a documented case of influenza virus infection associated with water exposure. Recreational water that has been treated at CDC recommended disinfectant levels does not likely pose a risk for transmission of influenza viruses. No research has been completed on the susceptibility of the H1N1 influenza virus to chlorine and other disinfectants used in swimming pools, spas, water parks, interactive fountains, and other treated recreational venues. However, recent studies have demonstrated that free chlorine levels recommended by CDC (1–3 parts per million [ppm or mg/L] for pools and 2–5 ppm for spas) are adequate to disinfect avian influenza A (H5N1) virus. It is likely that other influenza viruses such as novel H1N1 virus would also be similarly disinfected by chlorine.

Can H1N1 influenza virus be spread at recreational water venues outside of the water?

Yes, recreational water venues are no different than any other group setting. The spread of this novel H1N1 flu is thought to be happening in the same way that seasonal flu spreads. Flu viruses are spread mainly from person to person through coughing or sneezing of people with influenza. Sometimes people may become infected by touching something with flu viruses on it and then touching their mouth or nose.

Prevention & Treatment

What can I do to protect myself from getting sick?

There is no vaccine available right now to protect against this new H1N1 virus. There are everyday actions that can help prevent the spread of germs that cause respiratory illnesses like influenza.

Take these everyday steps to protect your health:

- Cover your nose and mouth with a tissue when you cough or sneeze. Throw the tissue in the trash after you use it.
- Wash your hands often with soap and water, especially after you cough or sneeze. Alcohol-based hand cleaners are also effective.
- Avoid touching your eyes, nose or mouth. Germs spread this way.
- Try to avoid close contact with sick people.
- Stay home if you are sick for 7 days after your symptoms begin or until you have been symptom-free for 24 hours, whichever is longer. This is to keep from infecting others and spreading the virus further.

Other important actions that you can take are:

- Follow public health advice regarding school closures, avoiding crowds and other social distancing measures.
- Be prepared in case you get sick and need to stay home for a week or so; a supply of over-the-counter medicines, alcohol-based hand rubs, tissues and other related items might be useful and help avoid the need to make trips out in public while you are sick and contagious.



What is the best way to keep from spreading the virus through coughing or sneezing?

If you are sick, limit your contact with other people as much as possible. If you are sick, stay home for 7 days after your symptoms begin or until you have been symptom-free for 24 hours, whichever is longer. Cover your mouth and nose with a tissue when coughing or sneezing. Put your used tissue in the waste basket. Then, clean your hands, and do so every time you cough or sneeze.

What is the best technique for washing my hands to avoid getting the flu?

Washing your hands often will help protect you from germs. Wash with soap and water or clean with alcohol-based hand cleaner. CDC recommends that when you wash your hands -- with soap and warm water -- that you wash for 15 to 20 seconds. When soap and water are not available, alcohol-based disposable hand wipes or gel sanitizers may be used. You can find them in most supermarkets and drugstores. If using gel, rub your hands until the gel is dry. The gel doesn't need water to work; the alcohol in it kills the germs on your hands.

What should I do if I get sick?

If you live in areas where people have been identified with new H1N1 flu and become ill with influenza-like symptoms, including fever, body aches, runny or stuffy nose, sore throat, nausea, or vomiting or diarrhea, you should stay home and avoid contact with other people. Staying at home means that you should not leave your home except to seek medical care. This means avoiding normal activities, including work, school, travel, shopping, social events, and public gatherings

If you have severe illness or you are at high risk for flu complications, contact your health care provider or seek medical care. Your health care provider will determine whether flu testing or treatment is needed

If you become ill and experience any of the following warning signs, seek emergency medical care.

In children, emergency warning signs that need urgent medical attention include:

- Fast breathing or trouble breathing
- Bluish or gray skin color
- Not drinking enough fluids
- Severe or persistent vomiting
- Not waking up or not interacting
- Being so irritable that the child does not want to be held
- Flu-like symptoms improve but then return with fever and worse cough

In adults, emergency warning signs that need urgent medical attention include:

- Difficulty breathing or shortness of breath
- Pain or pressure in the chest or abdomen
- Sudden dizziness
- Confusion
- Severe or persistent vomiting
- Flu-like symptoms improve but then return with fever and worse cough

Are there medicines to treat infection with this new virus?

Yes. CDC recommends the use of oseltamivir or zanamivir for the treatment and/or prevention of infection with the new H1N1 flu virus. Antiviral drugs are prescription medicines (pills, liquid or an inhaler) that fight against the flu by keeping flu viruses from reproducing in your body. If you get sick, antiviral drugs can make your illness milder and make you feel better faster. They may also prevent serious flu complications. During the current outbreak, the priority use for influenza antiviral drugs during is to treat severe influenza illness.

What is CDC's recommendation regarding "swine flu parties"?

"Swine flu parties" are gatherings during which people have close contact with a person who has novel H1N1 flu in order to become infected with the virus. The intent of these parties is to become infected with what for many people has been a mild disease, in the hope of having natural immunity to the novel H1N1 flu virus that might circulate later and cause more severe disease.

CDC does not recommend "swine flu parties" as a way to protect against novel H1N1 flu in the future. While the disease seen in the current novel H1N1 flu outbreak has been mild for many people, it has been severe and even fatal for others. There is no way to predict with certainty what the outcome will be for an individual or, equally important, for others to whom the intentionally infected person may spread the virus.

CDC recommends that people with novel H1N1 flu avoid contact with others as much as possible. They should stay home from work or school for 7 days after the onset of illness or until at least 24 hours after symptoms have resolved, whichever is longer.

Contamination & Cleaning

How long can influenza virus remain viable on objects (such as books and doorknobs)?

Studies have shown that influenza virus can survive on environmental surfaces and can infect a person for up to 2-8 hours after being deposited on the surface.

What kills influenza virus?

Influenza virus is destroyed by heat (167-212°F [75-100°C]). In addition, several chemical germicides, including chlorine, hydrogen peroxide, detergents (soap), iodophors (iodine-based antiseptics), and alcohols are effective against human influenza viruses if used in proper concentration for a sufficient length of time. For example, wipes or gels with alcohol in them can be used to clean hands. The gels should be rubbed into hands until they are dry.



What surfaces are most likely to be sources of contamination?

Germs can be spread when a person touches something that is contaminated with germs and then touches his or her eyes, nose, or mouth. Droplets from a cough or sneeze of an infected person move through the air. Germs can be spread when a person touches respiratory droplets from another person on a surface like a desk, for example, and then touches their own eyes, mouth or nose before washing their hands.

How should waste disposal be handled to prevent the spread of influenza virus?

To prevent the spread of influenza virus, it is recommended that tissues and other disposable items used by an infected person be thrown in the trash. Additionally, persons should wash their hands with soap and water after touching used tissues and similar waste.



What household cleaning should be done to prevent the spread of influenza virus?

To prevent the spread of influenza virus it is important to keep surfaces (especially bedside tables, surfaces in the bathroom, kitchen counters and toys for children) clean by wiping them down with a household disinfectant according to directions on the product label.

How should linens, eating utensils and dishes of persons infected with influenza virus be handled?

Linens, eating utensils, and dishes belonging to those who are sick do

not need to be cleaned separately, but importantly these items should not be shared without washing thoroughly first.

Linens (such as bed sheets and towels) should be washed by using household laundry soap and tumbled dry on a hot setting. Individuals should avoid “hugging” laundry prior to washing it to prevent contaminating themselves. Individuals should wash their hands with soap and water or alcohol-based hand rub immediately after handling dirty laundry.

Eating utensils should be washed either in a dishwasher or by hand with water and soap.

Response & Investigation

What is CDC doing in response to the outbreak?

CDC has implemented its emergency response. The agency’s goals are to reduce transmission and illness severity, and provide information to help health care providers, public health officials and the public address the challenges posed by the new virus. CDC continues to issue new interim guidance for clinicians and public health professionals. In addition, CDC’s Division of the Strategic National Stockpile (SNS) continues to send antiviral drugs, personal protective equipment, and respiratory protection devices to all 50 states and U.S. territories to help them respond to the outbreak.

What epidemiological investigations are taking place in response to the recent outbreak?

CDC works very closely with state and local officials in areas where human cases of new H1N1 flu infections have been identified. In California and Texas, where EpiAid teams have been deployed, many epidemiological activities are taking place or planned including:

- Active surveillance in the counties where infections in humans have been identified;
- Studies of health care workers who were exposed to patients infected with the virus to see if they became infected;
- Studies of households and other contacts of people who were confirmed to have been infected to see if they became infected;
- Study of a public high school where three confirmed human cases of H1N1 flu occurred to see if anyone became infected and how much contact they had with a confirmed case; and
- Study to see how long a person with the virus infection sheds the virus.

Who is in charge of medicine in the Strategic National Stockpile (SNS) once it is deployed?

Local health officials have full control of SNS medicine once supplies are deployed to a city, state, or territory. Federal, state, and local community planners are working together to ensure that SNS medicines will be delivered to the affected area as soon as possible. Many cities, states, and territories have already received SNS supplies. After CDC sends medicine to a state or city, control and distribution of the supply is at the discretion of that state or local health department. Most states and cities also have their own medicines that they can access to treat infected persons.

**Note: Much of the information in this document is based on studies and past experience with seasonal (human) influenza. CDC believes the information applies to the new H1N1 (swine) viruses as well, but studies on this virus are ongoing to learn more about its characteristics. This document will be updated as new information becomes available.*